LT3/18A/91/CT1

For treatment & control of Chronic Respiratory Diseases (CRD)

Tylosin 20% Powder

ASHTYL 20%

COMPOSITION: Each g contains:

Tylosin Tartrate BP (Vet.) equivalent to Tylosin . 20% w/w

PHARMACOLOGICAL CLASSIFICATION:

Antibiotics and Antibiotic combinations

Chicken & Turkeys: For the control of mortality caused by necrotic enteritis associated with Clostridium perfringens in broiler chickens. Chronic Respiratory Disease (CRD) caused by Mycoplasma gallisepticum, Mycoplasma synoviae.

Calves: Bovine pneumonia & calf scour

Pig: Swine dysentery & enzootic pneumonia.

DOSAGE AND ADMINISTRATION:

Chicken & Turkeys: 0.5 g of Tylosin tartrate per litre of drinking water or 1 teaspoonful of ASHTYL[™]20% in 2 litres of drinking water.

Calves: 0.5 g to 1 g Tylosin activity per calf administered orally twice daily for 7 -14 days. The product may be given in the milk or milk replacer.

Pigs: 3-6 mg Tylosin/kg b. wt. or 100 g Tylosin/ton of feed.

WITHDRAWAL PERIOD:

Broiler Chickens : 1 day Turkeys and Pigs : 0 day : 14 days

STORAGE: Store below 30°C in a dry place, protect from light.

CATEGORY OF DISTRIBUTION: Veterinary Medicines General Dealer (VMGD)

FOR DETAILED INFORMATION READ THE PACKAGE INSERT

KEEP OUT OF REACH OF CHILDREN

For treatment & control of Chronic Respiratory Diseases (CRD) Tylosin 20% Powder

ASHTYL[™]20% Tylosin 20% Powder

For treatment & control of Chronic Respiratory Diseases (CRD)



FOR VETERINARY USE ONLY

10 x 5 g

Mfg. Date: Date:

Batch No.:

Mfg. Lic. No.: KD-421

SAME SIZE DESIGN: 80 MM X 43 MM X 110 MM 10 X 5 GM CARTON

■ PANTONE 5473 C C M ● Y K ●







ASHTYL[™]20%

Tylosin 20% Powder

For treatment & control of Chronic Respiratory Diseases (CRD)